

DEPENDENCE OF GENERAL RELATIVISTIC ACCRETION ON BLACK HOLE SPIN

PARAMITA BARAI¹, TAPAS K. DAS², PAUL J. WIITA¹

¹ Department of Physics & Astronomy, Georgia State University, P.O. Box 4106, Atlanta GA 30302-4106
Email: barai@chara.gsu.edu, wiita@chara.gsu.edu

² N. Copernicus Astronomical Center, Bartycka 18, Warsaw 00-716 Poland
Email: tapas@camk.edu.pl

Draft version February 2, 2008

ABSTRACT

We propose a novel approach to the investigation of how black hole spin influences the behavior of astrophysical accretion. We examine the terminal behavior of general relativistic, multi-transonic, hydrodynamic advective accretion in the Kerr metric, by analyzing various dynamical as well as thermodynamic properties of accreting matter extremely close to the event horizon, as a function of Kerr parameter a . The examination of the transonic properties of such flows can be useful in self-consistently discriminating between prograde and retrograde relativistic accretion, and in studying the spectral signatures of black hole rotation.

Subject headings: accretion, accretion disks — black hole physics — hydrodynamics — relativity

1. INTRODUCTION

General relativistic (GR) accretion flows dive supersonically onto astrophysical black holes (BH) in order to satisfy the inner boundary condition imposed by the event horizon. Hence such accretion flows must be globally transonic (e.g., Fukue 1987; Chakrabarti 1989). This extremely hot, high density, ultra-fast flow can provide key features of the spectra of some galactic and extragalactic BH candidates. It is believed that most of the supermassive BHs powering active galactic nuclei (AGN) and stellar mass BHs in galactic microquasars possess non-zero values of the BH spin parameter, a , which might play a crucial role in determining their spectra and jet production (e.g., Rees 1984; Frank, King & Raine 1992; Kato, Fukue & Mineshige 1998). Hence it is very important to study the dynamical as well as the thermodynamic properties of GR transonic BH accretion very close to the event horizon and to investigate how BH spin modulates such behavior.

Study of the multi-transonic behavior of BH accretion was initiated by Abramowicz & Zurek (1981). Here we formulate and solve the conservation equations governing GR, multi-transonic, advective accretion in the Kerr metric to calculate the flow behavior extremely close ($\simeq 0.01$ gravitational radii) to the BH event horizon as a function of the Kerr parameter, a . Such a solution scheme for accretion onto a non-rotating BH has been outlined in Das (2004). We concentrate on accretion with sub-Keplerian angular momentum distributions. Such weakly rotating flows are exhibited in various physical situations, such as detached binary systems fed by accretion from OB stellar winds (Illarionov & Sunyaev 1975; Liang & Nolan 1984), semi-detached low-mass non-magnetic binaries (Bisikalo et al. 1998), and super-massive BHs fed by accretion from slowly rotating central stellar clusters (Illarionov 1988; Ho 1999 and references therein). Even for a standard Keplerian accretion disk, turbulence may produce such low angular momentum flow (e.g., Igumenshchev & Abramowicz 1999, and references therein). We find that our formalism is also valid for accretion with substantial angular momentum (as large as Keplerian) for retrograde flows.

2. FORMALISM

We consider non-self-gravitating, non-magnetized accretion. The gravitational radius is $r_g = GM_{BH}/c^2$ and we employ units where $G = c = M_{BH} = 1$. We use Boyer–Lindquist coordinates

with signature $-+++$, and an azimuthally Lorentz boosted orthonormal tetrad basis co-rotating with the accreting fluid. We define λ to be the specific angular momentum of the flow and neglect any gravo-magneto-viscous non-alignment between λ and a . We focus on the stationary axisymmetric solution of the following equations:

$$\nabla^\mu \mathfrak{Z}^{\mu\nu} = 0, \quad (\rho v^\mu)_{;\mu} = 0, \quad (1)$$

where $\mathfrak{Z}^{\mu\nu}$, v_μ , and ρ are the energy momentum tensor, the four velocity and the rest mass density of the accreting fluid, respectively, and the semicolon denotes the covariant derivative. The radial momentum balance condition may be obtained from $(v_\mu v_\nu + g_{\mu\nu}) \mathfrak{Z}^{\mu\nu} = 0$. Exact solutions of these conservation equations require knowledge of the accretion geometry and the introduction of a suitable equation of state. We concentrate on polytropic accretion for which $p = K\rho^\gamma$, where K and γ are the monotonic and continuous functions of the specific entropy density, and the constant adiabatic index of the flow, respectively. The specific proper flow enthalpy is taken to be $h = (\gamma - 1) \{ \gamma - (1 + a_s^2) \}^{-1}$, where a_s is the polytropic sound speed defined as $a_s = (\partial p / \partial \epsilon)_S^{1/2} = \Psi_1(T(r), \gamma) = \Psi_2(p, \rho, \gamma)$; here $T(r)$ is the local flow temperature, ϵ is the mass-energy density, and $\{\Psi_1, \Psi_2\}$ are known functions. The subscript S indicates that the derivative is taken at constant specific entropy. Hence, $a_s^2(r) = \gamma \kappa T(r) / (\mu m_H) = \Theta^2 T(r)$, where $\Theta = [\gamma \kappa / (\mu m_H)]^{1/2}$, μ is the mean molecular weight, m_H is the mass of the hydrogen atom, and κ is Boltzmann's constant.

We assume that the disk has a radius dependent local half-thickness, $H(r)$, and its central plane coincides with the equatorial plane of the BH. We use the vertically integrated model (Matsumoto et al. 1984) in describing black hole accretion disks where the equations of motion apply to the equatorial plane of the BH, assuming the flow to be in hydrostatic equilibrium in the transverse direction. Various dissipative processes are not explicitly taken into account. Even thirty years after the discovery of standard accretion disk theory (Shakura & Sunyaev 1973), modeling of viscous multi-transonic BH accretion, including proper heating and cooling mechanisms, is still quite an arduous task, even for Newtonian flows. The extremely large radial velocity close to the BH implies $\tau_{in} \ll \tau_{vis}$, with $\tau_{in}(\tau_{vis})$ the infall (viscous) timescales; hence our as-

sumption of inviscid flow is not unjustified, at least out to a few or to perhaps tens of r_g . One of the most significant effects of the introduction of viscosity would be the reduction of radial angular momentum. Hence study of the behavior of flow variables as functions of λ provides some insight about the flow behavior for viscous transonic accretion.

The temporal component of the first part of Eq. (1) leads to the conservation of specific flow energy $\mathcal{E}$ along each streamline as, $\mathcal{E} = hv_t$ (Anderson 1989). The Kerr metric in the equatorial plane of the black hole may be written as (e.g., Novikov & Thorne 1973)

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = -\frac{r^2 \Delta}{A} dt^2 + \frac{A}{r^2} (d\phi - \omega dt)^2 + \frac{r^2}{\Delta} dr^2 + dz^2, \quad (2)$$

where $\Delta = r^2 - 2r + a^2$, $A = r^4 + r^2 a^2 + 2ra^2$, and $\omega = 2ar/A$. The angular velocity Ω is

$$\Omega = -\frac{(g_{t\phi} + \lambda g_{tt})}{(g_{\phi\phi} + \lambda g_{t\phi})} = \frac{\frac{4a}{r} - \frac{\lambda(4a^2 - r^2 \Delta)}{A}}{\frac{A}{r^2} - \frac{4\lambda a}{r}}. \quad (3)$$

The normalization relation $v_\mu v^\mu = -1$, along with the above value of Ω , gives us the expression

$$v_t = \left[\frac{Ar^2 \Delta}{(1-u^2) \{A^2 - 8\lambda arA + \lambda^2 r^2 (4a^2 - r^2 \Delta)\}} \right]^{\frac{1}{2}}, \quad (4)$$

where u is the radial three velocity in the co-rotating fluid frame. Hence we obtain the conserved specific energy (which includes the rest mass energy) to be $\mathcal{E} = (\gamma - 1) [\gamma - (1 + \Theta^2 T)]^{-1} v_t$.

Note that our final calculations will provide the value of u and its related quantities rather than those of v_t . Hereafter, it is implied that any appearance of v_t in any mathematical expression is transformed in terms of u using Eq. (4).

We follow Abramowicz, Lanza & Percival (1997) to derive the expression for disk height $H(r)$ in our flow geometry because the corresponding equations in their calculations remain non-singular on the horizon and can accommodate a thin disk geometry as well as quasi-spherical flow. We obtain:

$$H(r) = \sqrt{2} r^2 \Theta T^{\frac{1}{2}} \left[\left(\frac{\gamma-1}{\gamma} \right) \left(\frac{1}{\gamma - (1 + \Theta^2 T)} \right) \left(\frac{1}{\psi} \right) \right]^{\frac{1}{2}}, \quad (5)$$

where $\psi = \lambda^2 v_t^2 - a^2 (v_t - 1)$. We define the quasi-invariant entropy accretion rate, $\dot{\Xi}$, as a quasi-constant multiple of the mass accretion rate, $\dot{M} = K^{\frac{1}{\gamma-1}} \dot{M}_{in}$, where $d\dot{\Xi}/dr = 0$ along a streamline only for a shock-free non-dissipative flow, and not for a flow containing shocks. The mass accretion rate ($\dot{M}$) and the entropy accretion rate are found to be

$$\dot{M} = 4\pi \rho \Theta T^{\frac{1}{2}} M r^2 \left[\frac{2\Delta(\gamma-1)\Theta^2 T}{\gamma(1-\Theta^2 T M^2) \{ \gamma - (1 + \Theta^2 T) \} \psi} \right]^{\frac{1}{2}}; \quad (6)$$

$$\dot{\Xi} = 4\sqrt{2} \pi r^2 u \sqrt{\frac{\Delta}{(1-u^2) \psi}} \left[\frac{a_s^2 (\gamma-1)}{\gamma \{ \gamma - (1 + a_s^2) \}} \right]^{\frac{1}{2} \left(\frac{\gamma+1}{\gamma-1} \right)}. \quad (7)$$

We simultaneously solve eqs. (4–7) along with the equation defining $\mathcal{E}$, to find the dynamical three-velocity gradient as

$$\frac{du}{dr} = \frac{\frac{2a_s^2}{(\gamma+1)} \left[\frac{r-1}{\Delta} + \frac{2}{r} - \frac{v_t \sigma \chi}{4\psi} \right] - \frac{\chi}{2}}{\frac{u}{(1-u^2)} - \frac{2a_s^2}{(\gamma+1)(1-u^2)u} \left[1 - \frac{u^2 v_t \sigma}{2\psi} \right]}, \quad (8)$$

where

$$\sigma = 2\lambda^2 v_t - a^2, \quad \chi = \frac{1}{\Delta} \frac{d\Delta}{dr} + \frac{\lambda}{(1-\Omega\lambda)} \frac{d\Omega}{dr} - \frac{\left(\frac{dg_{\phi\phi}}{dr} + \lambda \frac{dg_{t\phi}}{dr} \right)}{(g_{\phi\phi} + \lambda g_{t\phi})}. \quad (9)$$

Hereafter, we use the notation $[\mathcal{P}_3]$ for a set of values of $\{\mathcal{E}, \lambda, \gamma\}$ for a fixed Kerr parameter a (including the counter-rotating flows where a and λ have opposite signs), and $[\mathcal{P}_4]$ for a set of values of $\{\mathcal{E}, \lambda, \gamma, a\}$. By simultaneously setting the numerator and denominator of Eq. (8) to zero we find the sonic point conditions:

$$a_s|_c = \left[\frac{u^2 (\gamma+1) \psi}{2\psi - u^2 v_t \sigma} \right]_c^{\frac{1}{2}}, \quad u_c = \left[\frac{\chi \Delta r}{2r(r-1) + 4\Delta} \right]_c^{\frac{1}{2}}, \quad (10)$$

where the subscript c indicates that the quantities are to be measured at the critical (sonic) point(s). For a fixed $[\mathcal{P}_3]$ and a , we substitute the values of u_c and $a_s|_c$ in terms of r_c (Eq. 10) in the expression for $\mathcal{E}$, and obtain a polynomial in r_c , the solution of which provides the location of the sonic point(s) r_c . To determine the behavior of the solution near the sonic point, one needs to evaluate the value of du/dr at that point (the ‘critical velocity gradient’ $(du/dr)_c$) by applying L’Hospital’s rule to Eq. (8). After lengthy algebraic manipulations, we obtain the following quadratic equation which can be solved to obtain $(du/dr)_c$

$$\alpha \left(\frac{du}{dr} \right)_c^2 + \beta \left(\frac{du}{dr} \right)_c + \zeta = 0, \quad (11)$$

where the coefficients are

$$\begin{aligned} \alpha &= \frac{(1+u^2)}{(1-u^2)^2} - \frac{2\delta_1 \delta_5}{\gamma+1}, \quad \beta = \frac{2\delta_1 \delta_6}{\gamma+1} + \tau_6, \quad \zeta = -\tau_5; \\ \delta_1 &= \frac{a_s^2 (1-\delta_2)}{u(1-u^2)}, \quad \delta_2 = \frac{u^2 v_t \sigma}{2\psi}, \quad \delta_3 = \frac{1}{v_t} + \frac{2\lambda^2}{\sigma} - \frac{\sigma}{\psi}, \\ \delta_4 &= \delta_2 \left[\frac{2}{u} + \frac{uv_t \delta_3}{1-u^2} \right], \quad \delta_5 = \frac{3u^2 - 1}{u(1-u^2)} - \frac{\delta_4}{1-\delta_2} - \frac{u(\gamma-1-a_s^2)}{a_s^2 (1-u^2)}, \\ \delta_6 &= \frac{(\gamma-1-a_s^2) \chi}{2a_s^2} + \frac{\delta_2 \delta_3 \chi v_t}{2(1-\delta_2)}, \\ \tau_1 &= \frac{r-1}{\Delta} + \frac{2}{r} - \frac{\sigma v_t \chi}{4\psi}, \quad \tau_2 = \frac{(4\lambda^2 v_t - a^2) \psi - v_t \sigma^2}{\sigma \psi}, \\ \tau_3 &= \frac{\sigma \tau_2 \chi}{4\psi}, \quad \tau_4 = \frac{1}{\Delta} - \frac{2(r-1)^2}{\Delta^2} - \frac{2}{r^2} - \frac{v_t \sigma}{4\psi} \frac{d\chi}{dr}, \\ \tau_5 &= \frac{2}{\gamma+1} \left[a_s^2 \tau_4 - \{ (\gamma-1-a_s^2) \tau_1 + v_t a_s^2 \tau_3 \} \frac{\chi}{2} \right] - \frac{1}{2} \frac{d\chi}{dr}, \\ \tau_6 &= \frac{2v_t u}{(\gamma+1)(1-u^2)} \left[\frac{\tau_1}{v_t} (\gamma-1-a_s^2) + a_s^2 \tau_3 \right]. \end{aligned} \quad (12)$$

Note that all the above quantities are evaluated at r_c .




FIG. 1.— A typical flow topology for multi-transonic BH accretion: local Mach number plotted against distance from the event horizon of the BH, for $a = 0.2$, $\mathcal{E} = 1.0003$, $\lambda = 3.1$, and $\gamma = 4/3$. The curve **ABCD** represents accretion passing through the outer sonic point **B** ($r_o = 1094$), while **EBI** represents the self-wind. Flow along **GFH** passes through the inner sonic point **F** ($r_i = 3.796$) and encompasses a middle sonic point r_m ($r_m = 14.78$) denoted by an asterisk. A multi-transonic accretion flow passes through **AB**, then undergoes a shock which causes it to drop to the lower branch of the **GFH** curve.




FIG. 2.— Variation of terminal behavior of accretion flow variables, M_δ , T_δ , ρ_δ , and p_δ as functions of BH spin, a . The figure is drawn for a shock-free prograde flow passing through the outer sonic point, with $\mathcal{E} = 1.00001$; the curves (from left to right) correspond to values of $\lambda = 2.60, 2.12, 1.87, 1.55, 1.27, 1.04, 0.829$, and 0.609 . See text for details.

3. RESULTS

For a range of $[\mathcal{P}_4]$, one obtains three different values of r_c for a fixed $[\mathcal{P}_4]$, among which the largest and the smallest values correspond to the *X* type outer, r_o , and inner, r_i , sonic points, respectively. The *O* type middle sonic point, r_m , which is unphysical in the sense that no steady transonic solu-

tion passes through it, lies between r_i and r_o . For $\Xi(r_i) > \Xi(r_o)$ and $\Xi(r_i) < \Xi(r_o)$, one obtains multi-transonic accretion and a wind, respectively. If the accretion through r_o can be perturbed in a way so that it produces an amount of entropy exactly equal to $[\Xi(r_i) - \Xi(r_o)]$, supersonic flow through r_o can join the subsonic flow through r_i by developing a standing shock. The exact location of such a shock, as well as the details of the post-shock flow, may be obtained by formulating and solving the general relativistic Rankine-Hugoniot condition (Taub 1948; Thorne 1973) for our flow geometry. It is worth noting that the idea of shock formation in BH accretion is contested by some authors (see Narayan et al. 1998 and references therein for a review). However, the problem of not finding shocks in such works perhaps lies in the fact that only one (inner) sonic point close to the BH is explored by shock-free ADAF solutions.

For any $[\mathcal{P}_4]$, after obtaining $[r_i, r_m, r_o]$, we calculate the values of dynamical and acoustic velocities (Eq. 10) and the velocity gradient (Eq. 11) for $[r_i, r_o]$. These values are evaluated for a specific value of $\dot{M}$ ($= 1.0 \dot{M}_{Edd}$, in the results presented here; however, $\dot{M}$ could be chosen greater or less than the Eddington rate), and we can then compute $u(r)$, $a_s(r)$, the local Mach number, $M(r)$, and $T(r)$. Here we define $M(r) = u(r)/a_s(r)$, but because of our assumption of vertical equilibrium in an integrated disk model, $M(r_c) < 1$ (Matsumoto et al. 1984; Chakrabarti 1989). We compute the above quantities, the local vertically integrated density, $\Sigma(r)$, and pressure, $W(r)$ (Matsumoto et al. 1984), and other related dynamical or thermodynamic quantities by integrating the flow equations, from r_o and r_i down to the immediate vicinity of the event horizon, using a fourth order Runge-Kutta method. The flow could also be integrated for $r > r_c$ to study the subsonic part of the flow in detail. Figure 1 shows the integral curves of motion for a particular case. Similar topologies are obtained for any $[\mathcal{P}_4]$ allowing multi-transonic accretion flows. We find that $r_{i,pro} < r_{i,retro}$ for all $[\mathcal{P}_3]$ for which valid common solutions are available for both a and $-a$; however, for $a \gtrsim 0.2$, no such common $[\mathcal{P}_3]$ solutions can be found.

We define the “terminal” value of any accretion variable A at a radial distance $r_\delta = (r_e + \delta)$ as A_δ , where r_e is the event horizon, and $0 < \delta \ll 1.0$. Here we calculate A_δ by integrating the flow from r_c down to r_δ , so is clear that the dependence of A_δ on a can be studied at *any* r . For $\delta = 0.01$ and $M_{BH} = 10 M_\odot$, Fig. 2 shows the variations of M_δ , T_δ , ρ_δ and p_δ with BH spin for shock free prograde flows passing through r_o .

Note that as a flow through r_i does not connect r_e with infinity (such a flow folds back onto itself, see Fig. 1), solutions through r_i do not have *independent* physical existence, and can only be accessed if the supersonic flow through r_o undergoes a shock, so that it generates extra entropy, becomes subsonic, and produces the physical segment of the solution through r_i . If a shock forms, the calculation of A_δ for shocked flows boils down to the calculation of A_δ for flow through r_i . We find that for flows through r_i , the general profiles for the variations of A_δ with a remain exactly the same as that for flow through r_o ; only the magnitudes of various A_δ change. We obtain $M_\delta(r_i) < M_\delta(r_o)$ and $[T_\delta, \rho_\delta, p_\delta](r_i) > [T_\delta, \rho_\delta, p_\delta](r_o)$, because the shock compresses the flow and makes it hotter. Fig. 2 is drawn for $\gamma = 1.43$, but a similar figure could be drawn for any value of γ allowing multi-transonic accretion. We find that for both prograde and retrograde flows, $[T_\delta, \rho_\delta, p_\delta]$ non-linearly and monotonically correlate with γ while M_δ anti-correlates with γ . Multi-transonic

accretion does not exist for the same $[\mathcal{P}_3]$ for all values of a for prograde flows. Each curve segment in Fig. 2 shows the range of a for which a set of $\{\mathcal{E}, \lambda\}$ overlaps to produce three sonic points. Specifically, $\mathcal{E}$ is constant in Fig. 2 and each curve corresponds to a value of λ which allows multi-transonicity for that range of a . The allowed overlap range in a decreases as a increases.

For retrograde flows, the situation is different. For many combinations of $[\mathcal{P}_3]$, one obtains a large range of a (sometimes even covering the *complete* range of BH spin) for which one can have a *fixed* value of $\{\mathcal{E}, \lambda, \gamma\}$ producing multi-transonicity. Hence the variations of A_δ 's with a produce continuous curves covering almost the whole range $-1 < a \leq 0$. *Therefore multi-transonicity is more common for counter-rotating BH accretion.* We also found that, unlike prograde flows, retrograde flows with values of angular momentum even as high as Keplerian can demonstrate multi-transonic behavior.

From Fig. 2 we see that the correlations (of temperature, density and pressure) and the anti-correlation (of Mach number), with BH spin are non-linear but monotonic. Hence, rapidly rotating BHs will produce a hotter and denser *prograde* flow near the horizon. However, we again found that this is not true for counter-rotating flows, where the above-mentioned variations are not monotonic; rather they show distinct points of inflection at some intermediate value of a . The physical reason for such a behavior for retrograde flows is not clear to us. If one keeps $\{\mathcal{E}, \lambda, |a|\}$ fixed but decreases λ (which effectively mimics the introduction of viscosity to the system), $[T_\delta, \rho_\delta, p_\delta]$ non-linearly decreases and M_δ increases for prograde flow, whereas the trends are just the opposite for retrograde flow. These trends indicate how our results would be modified if our formalism were to be applied to study viscous transonic flow. Hence the study of A_δ as a function of a also opens up the possibility of distinguishing between the prograde and retrograde flows if the transonic properties of such flows can be measured.

4. DISCUSSION

Our formalism clearly allows the study of the dependence of the behavior of multi-transonic accretion on BH spin at any accretion length scale (including the radial shock location). Thus our work is important for understanding how BH spin influences several accretion related astrophysical phenomena, as only multi-transonic flows can produce shock waves in BH accretion disks (Das 2002; Das, Pendharkar & Mitra 2003, and references therein). Such hot and dense post-shock flow is important for understanding astrophysical phenomena such as spectral properties of BH candidates (Chakrabarti & Titarchuk 1995, and references therein), the formation and dynamics of accretion powered cosmic jets (Das & Chakrabarti 1999, and references therein; Das, Rao & Vadawale 2003), and the origin of QPOs in galactic sources (Das 2003, and references therein). Once we compute the exact shock location and related post-shock quantities (Barai, Das & Wiita, in preparation) for flow geometries presented in this Letter, it becomes straightforward to show how a determines the exact macrophysics of the post-shock flow right at the shock location, which in turn, may help determine how the properties of galactic and extragalactic jets and QPOs depend on the rotation of the BH.

One of the most tantalizing issues in relativistic astrophysics is to discover whether the spin of a BH can be determined from observations. By far the most discussed and accepted approach to this problem is through the study of the skewed shapes of fluorescent iron lines (see Reynolds & Nowak 2003 for a review). Our work provides an independent approach to this key question for hot, low angular momentum, prograde accretion flows, as well as for retrograde flows with substantial angular momentum.

PJW is grateful for continuing hospitality at the Department of Astrophysical Sciences at Princeton University. PB and PJW are partially supported by RPE funds to PEGA at GSU. TKD is supported by grants from the KBN.

REFERENCES

- Abramowicz, M. A., & Zurek, W. H. 1981, ApJ, 246, 314
 Abramowicz, M. A., Lanza, A., & Percival, M. J. 1997, ApJ, 479, 179
 Anderson, M. 1989, MNRAS, 239, 19
 Bisikalo, A. A., Boyarchuk, V. M., Chechetkin, V. M., Kuznetsov, O. A., & Molteni, D. 1998, MNRAS, 300, 39
 Chakrabarti, S. K. 1989, ApJ, 347, 365
 Chakrabarti, S. K., & Titarchuk, L. G. 1995 ApJ, 455, 623
 Das, T. K. 2002, ApJ, 577, 880
 Das, T. K. 2003, ApJ, 588, L89
 Das, T. K. 2004, MNRAS, 375, 384
 Das, T. K., & Chakrabarti, S. K. 1999, Class. Quantum Grav. 16, 3879
 Das, T. K., Pendharkar, J. K., & Mitra, S. 2003, ApJ, 592, 1078
 Das, T. K., Rao, A. R., & Vadawale, S. R. 2003, MNRAS, 343, 443
 Frank, J., King, A. R., & Raine, D. J. 1992, Accretion Power in Astrophysics (2nd ed.; Cambridge: Cambridge Univ. Press)
 Fukue, J. 1987, PASJ, 39, 309
 Ho, L. C. 1999, in Observational Evidence For Black Holes in the Universe, ed. S. K. Chakrabarti (Dordrecht: Kluwer), 153
 Igumenshchev, I. V., & Abramowicz, M. A. 1999, MNRAS, 303, 309
 Illarionov, A. F. 1988, Soviet Astron., 31, 618
 Illarionov, A. F., & Sunyaev, R. A. 1975, A&A, 39, 205
 Kato, S., Fukue, J., & Mineshige, S., 1998, Black-hole Accretion Disks. (Kyoto: Kyoto University Press)
 Liang, E. P. T., & Nolan, P. L. 1984, Space Sci. Rev., 38, 353
 Matsumoto, R., Kato, S., Fukue, J., & Okazaki, A. T. 1984, PASJ, 36, 71
 Narayan, R., Mahadevan, R., & Quataert, E. 1998, in Theory of Black Hole Accretion Disks, ed. M. A. Abramowicz, G. Björnsson, J. E. Pringle (Cambridge: Cambridge University Press), 148
 Novikov, I., & Thorne, K. S. 1973, in Black Holes, ed. C. De Witt, B. De Witt (New York: Gordon and Breach), 343
 Rees, M. J. 1984, ARA&A, 22, 471
 Reynolds, C. S., & Nowak, M. A. 2003, Phys. Rep., 377, 389
 Shakura, N. I., & Sunyaev, R. A. 1973, A&A, 24, 337
 Taub, A. H. 1948, Phys. Rev. 74, 328
 Thorne, K. S. 1973, ApJ, 179, 897